

See discussions, stats, and author profiles for this publication at: <http://www.researchgate.net/publication/282543966>

Larval diet alters larval growth rates and post-metamorphic performance in the marine gastropod *Crepidula fornicata*

ARTICLE *in* MARINE BIOLOGY · JULY 2015

Impact Factor: 2.39 · DOI: 10.1007/s00227-015-2696-7

READS

21

2 AUTHORS, INCLUDING:



Jan Pechenik

Tufts University

111 PUBLICATIONS 3,737 CITATIONS

SEE PROFILE

Larval diet alters larval growth rates and post-metamorphic performance in the marine gastropod *Crepidula fornicata*

Jan A. Pechenik¹ · Abigail S. Tyrell²

Received: 18 March 2015 / Accepted: 2 July 2015 / Published online: 19 July 2015
© Springer-Verlag Berlin Heidelberg 2015

Abstract Some larval experiences can produce “latent effects” on post-metamorphic growth or survival. While it is known that periods of starvation during larval development can cause such latent effects, the effect of larval diet on post-metamorphic growth has not been studied. As global climate change and ocean acidification are expected to decrease phytoplankton concentrations and alter both phytoplankton species composition and nutritional characteristics, we examined the impact of 3 phytoplankton species (*Isochrysis galbana*, clone T-ISO; *Pavlova lutheri*, clone MONO; and *Dunaliella tertiolecta*, clone DUN) on larval growth and subsequent post-metamorphic fitness in the slipper snail *Crepidula fornicata*. Once larvae metamorphosed, the juveniles were all reared on the diet that produced the fastest growth, T-ISO, to look for latent effects of larval diet on juvenile growth. In all experiments, larvae grew most quickly on T-ISO; diet did not affect relative rates of shell and tissue growth. In 2 of the 4 experiments conducted on the effects of diet quality, larvae reared on T-ISO metamorphosed into juveniles that grew significantly faster than those that had been raised on the other phytoplankton species, indicating clear latent effects of dietary experience and suggesting parent-related genetic variation in susceptibility to this type of stress. Rearing larvae at a

very low food concentration of T-ISO (1×10^4 cells ml⁻¹) until metamorphosis also produced severe latent effects on juvenile growth, reducing juvenile growth rates by more than 30 %. These data provide yet another example of how stresses experienced during larval development can influence post-metamorphic performance, and add another level of complexity to attempts at predicting the future consequences of environmental change on marine community structure and species interactions.

Introduction

The microscopic, planktonic larval stages of many benthic marine invertebrates must feed on phytoplankton for days or weeks before becoming competent to metamorphose (reviewed by Thorson 1950; Crisp 1974; Morgan 1995). The effects of food concentration and diet on larval growth rates and time to metamorphic competence have been studied for a number of marine invertebrate species (Pechenik 1987; His and Seaman 1992; Hansen 1993; Anger 1995; Basch 1996; Klinzing and Pechenik 2000; McEdward and Qian 2001), in part because the extent to which diet slows larval growth and prolongs larval life should increase the extent of planktonic mortality (Rumrill 1990; Morgan 1995; Vaughn and Allen 2010; Byrne 2011) and dispersal away from the parental habitat (Pechenik 1999; O’Connor et al. 2007; Cowen and Sponaugle 2009). However, inadequate food concentrations during larval development can also have detrimental effects following metamorphosis, so-called “latent effects”: post-metamorphic consequences of stresses experienced during embryonic or larval development (reviewed by Pechenik 2006). For example, juveniles of the marine gastropod *Crepidula fornicata* reared under ideal conditions typically grew at least 30 % more slowly for at least the

Communicated by G. Chapman.

Reviewed by undisclosed experts.

✉ Jan A. Pechenik
jan.pechenik@tufts.edu

¹ Biology Department, Tufts University Medford, Medford, MA 02155, USA

² The School of Marine and Atmospheric Sciences, Stony Brook University, Stony Brook, NY 11794, USA

first several days after metamorphosis if the larvae had been starved earlier in development for even just 2 days (Pechenik et al. 1996a, b, 2002). Similarly, larvae of the related species *C. onyx* grew significantly slower as juveniles if they had been reared at a low food concentration as larvae (Chiu et al. 2007), as did the mussel *Mytilus galloprovincialis* (Phillips 2002). To the extent that there is an escape in size from predation, competition, or bulldozing following metamorphosis (Gosselin and Qian 1997; Hunt and Scheibling 1997; Pechenik et al. 2010; Jennings and Hunt 2011), such reduced juvenile growth rates would likely increase the extent of early post-metamorphic mortality.

Larval growth rates can also vary substantially with the species of phytoplankton provided as food (Pechenik and Fisher 1979; Klinzing and Pechenik 2000; reviewed by Pechenik 1987). In particular, in previous studies, the larvae of *C. fornicata* grew much more rapidly on a diet of *Isochrysis galbana* (clone T-ISO) than on either *Pavlova lutheri* (clone MONO) or *Dunaliella tertiolecta* (clone DUN; Klinzing and Pechenik 2000), and grew more rapidly on a diet of *D. tertiolecta* than on a diet of *P. lutheri* (Lucas and Costlow 1979). The impact of different larval diets on post-metamorphic performance has not previously been reported. Such information is becoming increasingly relevant, however, as climate change and ocean acidification have the potential to alter phytoplankton species ranges and cause shifts in both species composition (Hinga 2002; Tortell et al. 2002; Hayes et al. 2005; Kim et al. 2006; Leu et al. 2013) and the nutritional content of individual microalgal species (Hoogstraten and Timmermans 2012; Rossoll et al. 2012; Wynne-Edwards et al. 2014).

In this study, we sought to determine whether phytoplankton species that alter larval growth rates in the marine gastropod *Crepidula fornicata*, based on results from previous studies with this species (Lucas and Costlow 1979; Klinzing and Pechenik 2000), can also affect subsequent post-metamorphic growth rates on an ideal diet. We also asked whether diet can alter not only rates of larval shell growth, but also relative rates of shell and tissue growth. Finally, we examined the impact on juvenile development of rearing larvae on an ideal diet (T-ISO) at a low phytoplankton concentration. Two prior studies with *C. fornicata* (Pechenik et al. 1996a, 2002) found no latent effects on juvenile development after feeding larvae at a low food concentration (1×10^4 cells ml^{-1}) of T-ISO for 2 or 4 days; in the present study we reared larvae at that same low-food concentration from hatching until metamorphosis (typically ~10–15 days). Such treatment did produce latent effects in the related species *C. onyx* (Chiu et al. 2007). In our study, we also examined the effects of this treatment on relative rates of shell and tissue growth during larval development.

Crepidula fornicata is ideal for such studies: larvae hatch at shell lengths of about 450 μm and grow quickly

(typically 50–100 $\mu\text{m day}^{-1}$; Pechenik et al. 1996c; Hilbish et al. 1999; Klinzing and Pechenik 2000), making them easy to measure non-destructively (Pechenik et al. 2002); larvae typically become competent to metamorphose within 2 weeks at room temperature at large sizes, at shell lengths approximately 850–1200 μm (Pechenik and Heyman 1987; Pechenik et al. 1996c); larval shell length and organic tissue weight are linearly related for this species under normal rearing conditions (Pechenik 1980), so that shell growth is an accurate measure of tissue growth; competent larvae can be triggered to metamorphose within 6–8 h, simply by elevating ambient K^+ concentration by 15–20 mM (Pechenik and Gee 1993); and both larvae and juveniles can be reared in the laboratory with negligible mortality, so that any latent effects encountered cannot be due to selective mortality for particular genotypes (Pechenik et al. 1996c, 2002; Klinzing and Pechenik 2000). Finally, although *C. fornicata* is native to the East Coast of the United States, it is now widely distributed among coastal communities in many parts of the world (Blanchard 1997; Bohn et al. 2012), making it potentially useful as a biomonitor of climate change.

Materials and methods

General procedures

Several stacks of adults were collected intertidally at Nahant, Massachusetts (collecting permit issued by the Commonwealth of Massachusetts, Division of Marine Fisheries) in September 2013 (Experiments 1 and 2), early November 2013 (Experiment 3), March 2014 (Experiment 4), and May 2014 (Experiment 5), and brought to the laboratory where they were maintained on a mixed diet of *Isochrysis galbana* (clone T-ISO) and *Dunaliella tertiolecta* (clone DUN; Pechenik et al. 1996a, b) until larvae were released 3–10 days later. Larvae were then retained on a 150 μm Nitex mesh filter and transferred to seawater that had been filtered to 0.45 μm (Pechenik et al. 1996a).

Five experiments were conducted. The first 4 experiments were designed to detect latent effects resulting from rearing larvae on different algal diets: larvae were raised on a diet of either T-ISO (*Isochrysis galbana*), MONO (*Pavlova lutheri*), or DUN (*Dunaliella tertiolecta*); cell sizes range from about 3–5 μm for T-ISO to about 8–10 μm for DUN (Pechenik and Fisher 1979). All 3 species have been widely used for molluscan aquaculture (e.g., Volkman et al. 1989) and their effects on the survival and growth of *C. fornicata* larvae have been examined in previous studies (e.g., Lucas and Costlow 1979; Klinzing and Pechenik 2000) although latent effects on post-metamorphic development have not been previously examined. Phytoplankton

were cultured at 18 °C using f/2 medium and standard techniques (Guillard and Ryther 1962), and harvested at concentrations of $\sim 1\text{--}3 \times 10^6$ cells ml^{-1} . Larvae (14–19 per dish) were reared in 45 ml of 0.45 μm filtered seawater (salinity 30) at room temperature (~ 23 °C), with phytoplankton at $\sim 18 \times 10^4$ cells ml^{-1} as in previous studies with this species (Klinzing and Pechenik 2000; Pechenik et al. 1996a, b, c; Pires et al. 2000). Five dishes of larvae were established for each treatment. Three dishes of larvae were used to monitor larval growth and to obtain juveniles after metamorphosis. One additional dish of larvae was used for tissue and shell weight analyses, to determine if rates of shell and tissue growth were affected by diet to the same degree, while the fifth dish of larvae was used to monitor the onset of metamorphic competence. Seawater and food were changed every 2–3 days; in spot checks, phytoplankton concentrations in Experiments 1–4 never fell below $\sim 10 \times 10^4$ cells ml^{-1} , well above the critical concentration below which ingestion rates are impacted ($\sim 4 \times 10^4$ cells ml^{-1} , Pechenik and Eyster 1989).

In a final experiment (Experiment 5), we reared larvae at high (18×10^4 cells ml^{-1}) or low (1×10^4 cells ml^{-1}) concentrations of the diet on which larvae grew the fastest (T-ISO) for the entire larval period, to determine whether low concentrations of a nutritious microalga produces latent effects on post-metamorphic growth in this species as it does in *C. onyx* (Chiu et al. 2007). The two phytoplankton concentrations were chosen based on the results of previous studies (Pechenik et al. 1996a; Klinzing and Pechenik 2000; Chiu et al. 2007).

For all experiments, treatments were begun on Day 0, 1–2 days after the larvae hatched, at which time mean initial shell length was determined for at least 12 larvae at $50\times$ or $63\times$ (Table 1), using a dissecting microscope equipped with an ocular micrometer; there is little within-brood variation in shell length at hatching for this species (Pechenik et al. 1996a, 2002) so that these measurements suffice for estimating initial larval size. Larvae were then measured at least 2 more times during development to determine growth rates. Final larval shell lengths were determined when larvae had reached an average size of ~ 850 μm , after different lengths of time depending on larval growth rates (e.g., in Experiment 1, after 9 days for larvae reared on T-ISO and after 13 days for larvae reared on MONO).

Within each experiment, once larvae reached shell lengths of ~ 850 μm , the larvae from one replicate dish were tested periodically for metamorphic competence by raising the concentration of KCl in seawater by 20 mM; competent larvae typically metamorphose in response to this treatment within 6–8 h (Pechenik and Gee 1993), and without any impact on post-metamorphic growth rates (Eyster and Pechenik 1988). If more than 50 % of the tested larvae metamorphosed within 6–8 h of exposure to

Table 1 Sizes of larvae and juveniles used for the shell and tissue weight determinations reported in Fig. 7

Experiment	N	Mean shell length (μm)	SD
Experiment 1			
Larvae			
T-ISO	44	835.3	160.3
MONO	43	792.6	120.5
Juveniles			
T-ISO	15	3097.6	870.5
MONO	12	3489.1	367.7
Experiment 2			
Larvae			
T-ISO	44	924.7	55.1
MONO	47	847.9	102.9
DUN-3	44	825.7	76.7
DUN-6	44	865.7	85.3
DUN-9	37	807.0	83.3
Juveniles			
T-ISO	13	4221.1	183.2
MONO	13	3012.8	489.8
DUN-3	12	3918.3	194.4
DUN-6	12	3827.9	288.4
DUN-9	12	3352.3	519.3
Experiment 3			
Larvae			
T-ISO	17	813.0	35.7
MONO	17	831.4	28.4
DUN-5	18	825.3	30.1
DUN-10	17	807.9	38.4
Experiment 4			
Larvae			
T-ISO	12	834.1	114.2
MONO	12	839.4	101.5
Experiment 5			
Larvae			
T-ISO high food	31	963.8	106.9
T-ISO low food	27	954.2	80.0

the inducer, metamorphosis was then induced in the other replicates of the same diet on the following day. Larvae that failed to metamorphose within the 6–8 h exposure periods (“pre-competent” larvae—Pechenik and Gee 1993) were preserved in 10 % formalin buffered to a pH of approximately 8.0 with sodium borate, for later determinations of shell and tissue weights. Larvae that had been reared expressly for determining shell and tissue weight were also preserved at this time.

Following metamorphosis, 12–15 juveniles haphazardly chosen from each treatment (up to 5 from each replicate) were reared individually in 45 ml of

filtered seawater (salinity 30, ~23 °C) on a diet of T-ISO at $\sim 18 \times 10^4$ cells ml⁻¹, which generally supports rapid juvenile growth (typically 100–200 $\mu\text{m day}^{-1}$) with negligible mortality (Pechenik et al. 1996b, c, 2002; Klinzing and Pechenik 2000); young juveniles of this species grow at constant rates above food concentrations of about 7×10^4 cells ml⁻¹ (Eyster and Pechenik 1988; Pechenik and Eyster 1989). Note that all juveniles were reared on T-ISO, regardless of what the larval diet had been. Water was changed every 1–2 days. On days that the water was not changed, juveniles were given enough additional T-ISO to restore the total cell concentration to approximately 18×10^4 cells ml⁻¹. In spot checks, cell concentrations never fell below 10×10^4 cells ml⁻¹.

Juvenile shells were measured at least twice (at 12–50 \times magnification, depending on their size): once within 24 h of metamorphosis and once again 3–8 days later.

Shell and tissue weights were determined as follows. Weights were determined for both larvae and juveniles in Experiments 1 and 2, but only for larvae in Experiments 3, 4, and 5. Preserved larvae and juveniles were quickly rinsed in deionized water to remove adhering salts and preservative and then dried in small pre-weighed aluminum foil cups at 50 °C for at least 48 h before being weighed to the nearest 0.001 mg on a Mettler Toledo balance, to determine total dry weight. For larval weight determinations in Experiments 1–4, we had 2–4 replicates per diet, each with 4–10 individuals per replicate; for Experiment 5, we had 3 replicates with 11–14 individuals per replicate. For juveniles, we used 3–9 replicates per treatment, with one individual per replicate. After weighing, samples were combusted at 500 °C for at least 12 h to determine organic (tissue) weight by quantifying weight loss following combustion (Paine 1964; Pechenik 1980; Pechenik and Eyster 1989). Control foil containers did not lose weight during this process. Shell (inorganic) weights were determined by subtracting the empty container weight from the container's post-combustion weight. Mean shell lengths of larvae and juveniles are given in Table 1.

Details of particular experiments are given below.

Experiments 1–4: Impact of larval diet on juvenile growth

In Experiment 1, larvae were raised on a diet of T-ISO, MONO, or DUN. Larval survival on DUN was very poor in this experiment (see Results); surviving larvae were transferred to a diet of T-ISO after 18 days, but only one of those larvae survived to metamorphosis so that juvenile growth rates could not be documented. Following the metamorphosis of larvae reared on the other 2 diets, 15 juveniles (5 from each replicate) that had been reared as larvae on

T-ISO and 15 juveniles that had been reared as larvae on MONO were reared on a diet of T-ISO for another 15 days, again at room temperature (23 °C). Juveniles were changed to new seawater and phytoplankton daily. Shells were measured at 12–50 \times , depending of their size, to determine juvenile shell growth rates. After making the final shell measurements, juveniles were preserved in 10 % buffered formalin as described above, for later analysis of shell and tissue weight.

For Experiments 2 and 3, larvae were raised on the same 3 diets used in Experiment 1, all at approximately 18×10^4 cells ml⁻¹, as in Experiment 1. Because larvae had survived poorly on a diet of DUN in the previous experiment, we raised groups of larvae on DUN in Experiment 2 for 3, 6, or 9 days and in Experiment 3 for 5 or 10 days, and then switched them to a diet of T-ISO for the rest of larval development. Larvae were measured a number of times during larval development.

Toward the end of larval rearing in Experiment 2 we noticed that our T-ISO phytoplankton culture had become mildly contaminated with DUN. Therefore we include data for only the first 4 days of juvenile development for juvenile growth rate determinations, during which time T-ISO cells still dominated the phytoplankton culture. Shell and tissue weights were determined when juveniles were 14–18 days old.

Experiment 4 was similar to the previous experiments except that larvae were reared on T-ISO for the entire larval period or on MONO for 3 (treatment MONO-3), 6, or 9 days, after which they were transferred to a diet of T-ISO for the duration of the study. For the T-ISO and MONO-3 data, larvae were measured on days 0, 3, and 6. For the MONO-6 and MONO-9 data, larvae were only measured twice, on days 0 and 6.

Experiment 5: Impact of rearing larvae at a low T-ISO food concentration

In this experiment, larvae were raised at two different concentrations of T-ISO (18×10^4 cells ml⁻¹ or 1×10^4 cells ml⁻¹) until metamorphosis, to determine whether rearing larvae of this species on a high-quality diet at a low food concentration produces latent effects on juvenile growth. Larvae reared at the high food concentration were measured on days 0, 3, and 9, while those reared at the low food concentration were measured on days 0, 4, and 10.

Statistical analyses

The effects of larval diet on mean larval shell growth rates, mean juvenile shell growth rates, mean larval tissue weight as a percentage of overall mass, and mean juvenile tissue

weight as a percentage of overall mass were analyzed using: (1) ANOVA followed by Tukey's multiple comparisons test if there were more than two groups being compared; (2) Student's *t* test if there were only two groups being compared and data for both groups were normally distributed with similar variances; (3) Welch's *t* test if the variances of the two groups being compared were unequal; or (4) Mann–Whitney U test if either of the two groups being compared had a non-normal distribution. There were no significant differences in mean larval growth rates among replicates (ANOVA, $P > 0.10$); growth rate data from replicates were therefore combined for further analysis. Data on percent larval survival were arc-sine transformed before analysis. Data were analyzed using GraphPad Prism 4.0.

Results

Experiment 1

Larval growth rates were linear in all treatments (T-ISO $R^2 = 0.973$; MONO $R^2 = 0.98$; DUN $R^2 = 0.884$). Larvae grew significantly faster on T-ISO than on either MONO or DUN, and grew significantly more slowly on DUN than on MONO ($F_{(3,131)} = 54.3$, $P < 0.0001$; Tukey's tests, $P < 0.05$; Fig. 1A). After 9 days of being reared on a diet of DUN, larval shell length plateaued at about 650 μm ; larvae grew even more slowly on this diet during days 9–20 (Tukey's test, $P < 0.05$; Fig. 1A), despite having been transferred to a diet of T-ISO after day 17. Mortality was 91 % by day 20 for larvae being reared on DUN, but was less




Fig. 1 Impact of larval diet on larval growth rates (A) and on subsequent juvenile growth rates (B, C). After metamorphosis, juveniles were reared on a diet of T-ISO. Larvae were reared on 1 of 3 diets, as indicated. For larvae reared on DUN, larval growth rates are shown for two different time periods (A). For larvae reared on T-ISO or

MONO, growth rates were based on mean initial size and sizes after 10–14 days of development. The number of individuals measured is indicated within each bar. Different letters above each bar represent significant differences between means (Tukey *t* tests, $P < 0.05$). For juveniles, 13–15 individuals were measured per treatment

than 7 % in any replicate for larvae in other treatments. Only a few larvae raised on DUN survived to the juvenile stage, so that juvenile growth rates could not be determined. Such poor performance on DUN was not seen in subsequent experiments (see below).

Larvae raised on MONO grew nearly 38 % more slowly than those raised on T-ISO (Fig. 1A). Despite this large difference in larval growth rates, differences in larval diet produced no significant differences in juvenile growth rates, either for the first 3 days after metamorphosis (Fig. 1B; $t = 0.04$, $df = 26$, $P = 0.97$) or for the first 15 days after metamorphosis (Fig. 1C; Mann–Whitney U test, $U = 79.0$, $P = 0.61$). Juvenile mortality ranged between 0 and 13 % ($N = 15$ individuals per treatment).

Experiment 2

Larval mortality was generally below 7 % in all treatments. Larvae grew at constant rates over time on all diets (R^2 for length versus time varied only from 0.93 to 0.99; Fig. 2) but growth rates differed on different diets (see below). Growth rates for larvae reared on DUN for only 3, 6, or 9 days never recovered to control levels even after larvae had been transferred to T-ISO for the remainder of the larval rearing period (Fig. 2).

Although our T-ISO algal culture became contaminated with DUN toward the end of larval rearing in this experiment, larvae reared on a diet of “T-ISO” exhibited higher mean growth rates over the course of the 15–19 days larval period than larvae reared in any of the other treatments, including those in which larvae were reared purely on DUN for up to 9 days (Figs. 2, 3A; $F_{(3,131)} = 54.3$, $P < 0.0001$;




Fig. 2 Effect of diet on larval growth in Experiment 2. Data are the means of measurements made on 37–46 larvae on each day. Error bars (SD) are shown only for one treatment, but were similar for other treatments

Tukey’s tests, $P < 0.05$). Larvae reared on MONO grew about 33 % more slowly than those reared on T-ISO, and about 18 % more slowly than those reared on DUN for either 3 or 6 days, but at equal rates to those reared on DUN for 9 days (Tukey’s tests, $P > 0.05$; Fig. 3A).

Over the first 4 days after metamorphosis, juveniles from larvae that had been raised on DUN for 9 days (treatment DUN-9) grew significantly more slowly than those that had been reared on T-ISO or MONO, or on DUN for only 6 days (DUN-6; $F_{(4,57)} = 5.13$, $P = 0.0013$; Tukey’s tests, $P < 0.05$; Fig. 3B). Indeed, those juveniles grew about 35 % more slowly than those that had been reared on T-ISO as larvae. There were no other significant differences in juvenile shell growth rates among treatments (Tukey’s tests, $P > 0.05$; Fig. 3B). Juvenile mortality over those first 4 days was 0 % for treatments T-ISO, MONO, and DUN-6; 7 % for DUN-3; and 20 % for DUN-9 ($N = 15$ individuals in each treatment).

Experiment 3

Larval mortality was less than 5 % in all treatments and growth was linear on all diets (DUN-10 $R^2 = 0.97$; $R^2 > 0.99$ for larvae in all other treatments). Larvae again grew substantially more quickly when reared on T-ISO than when reared on the other two diets ($F_{(6,128)} = 73.0$; $P = 0.006$); in particular, larvae reared on MONO grew about 40 % more slowly than those reared on T-ISO (Fig. 4A). Larvae also grew significantly more slowly on a diet of DUN than on T-ISO (Fig. 4A; Tukey’s test, $P < 0.05$), and growth rates for larvae reared on DUN for 10 days were as low as those for larvae reared on MONO (Tukey’s test, $P > 0.10$).

Juveniles that had been reared as larvae on MONO grew significantly more slowly than those that had been reared on T-ISO as larvae, for at least the first 5 days after metamorphosis (Tukey’s test, $P < 0.05$; Fig. 4B). Indeed, 5 individuals (of the 12 monitored) that had been reared on MONO as larvae grew at less than $32 \mu\text{m day}^{-1}$ as juveniles, on a diet of T-ISO. There were no significant differences in juvenile growth rates for individuals that had been reared as larvae on T-ISO or on DUN, even for those that had been reared on DUN for 10 days (Tukey’s test, $P > 0.10$), despite the difference in larval growth rates.

Experiment 4

Larval mortality was below 5 % in all treatments, and larvae grew at constant rates (T-ISO $R^2 = 0.993$; MONO-3 $R^2 = 0.990$; larvae were only measured at two different times for the other two treatments). As in all previous experiments, larvae grew significantly faster on a diet of T-ISO than on MONO (ANOVA $F_{(5,108)} = 42.9$,




Fig. 3 Effect of larval diet on larval (A) and juvenile (B) growth rates in Experiment 2. Larvae were reared on T-ISO or MONO for all of larval development; larvae were reared on DUN for only 3, 6, or 9 d, as indicated, and were then transferred to a diet of T-ISO for

the rest of their development. The number of individuals measured is indicated within each bar. Different letters above bars indicate means that differ significantly (Tukey’s tests, $P < 0.05$)




Fig. 4 Effect of larval diet on larval (A) and juvenile (B) growth rates in Experiment 3. Larvae were reared on T-ISO or MONO for all of development, but reared on DUN for either 5 d (DUN-5) or 10 d (DUN-10). Larvae were measured at hatching and after 5 or 8 d of larval development. All juveniles were reared on a diet of T-ISO and

measured on the days indicated. Larval growth rates were based on shell measurements for 17–21 larvae on each day. For juveniles (B), numbers within bars indicate the number of individuals monitored. Bars are means + one SD. Lower case letters represent individual treatments, referred to in the text of the Results section

$P < 0.0001$; Tukey tests, $P < 0.05$; Fig. 5A), even though individuals raised on MONO were transferred to a diet of T-ISO after 3, 6, or 9 days. Larvae grew significantly more slowly when reared on a diet of MONO for 6 or 9 days (treatments MONO-6, MONO-9) and then transferred to T-ISO than when reared on MONO for only 3 days (MONO-3) before their transfer to T-ISO (Tukey test,

$P < 0.05$). Nevertheless, juvenile growth rates were not affected by larval diet, regardless of the time period over which growth rates were measured (Fig. 5B). Note that although a significant P -value was obtained by ANOVA for juvenile growth rates over the first 4 days after metamorphosis (Fig. 5B), no paired comparisons differed significantly (Tukey’s tests, $P > 0.05$ for all comparisons).




Fig. 5 Effect of larval diet on larval (A) and juvenile (B) growth rates in Experiment 4. Larvae were reared on TISO for all of larval development, but reared on MONO for either 3, 6, or 9 d, as indicated, before being transferred to a diet of T-ISO for the rest of larval development. Larvae were measured at hatching and again on day

3 and 6 as indicated; data are means of 16–20 measurements + SD. For determinations of juvenile growth rates, 11–13 individuals were measured at each of the sampling periods, including the day after metamorphosis (Day 0). There were no significant differences among means for either group of data (Tukey's tests, $P < 0.05$)




Fig. 6 Effects of larval diet concentration on larval and juvenile growth rates of *C. formicata*. Larvae were reared at either 18×10^4 cells ml^{-1} T-ISO (high food) or 1×10^4 cells ml^{-1} T-ISO (low food). Larvae were measured at hatching and again after 3–4 days and after 9 days, after which metamorphosis was induced. After metamorphosis, all juveniles were reared at the high food concentration. *Significant differences between means ($P \leq 0.01$)

Experiment 5

Larvae grew significantly more slowly on a diet of T-ISO at 1×10^4 cells ml^{-1} than at 18×10^4 cells ml^{-1} (days

0–4, $t = 10.8$, $df = 80$, $P < 0.0001$; days 0–9, Mann–Whitney $U = 23.0$, $P < 0.0001$; Fig. 6). All larvae exhibited linear growth (high food $R^2 = 0.999$, low food $R^2 = 0.995$), and mortality was under 5 % in all replicates. Following metamorphosis, when all juveniles were reared at the high food concentration, juveniles that had been reared as larvae at the low food concentration showed latent effects for at least the first 6 days after metamorphosis, growing 34–43 % more slowly than individuals that had been raised as larvae at the high food concentration (days 0–3, Mann–Whitney $U = 37.0$, $P = 0.01$; days 0–6, $t = 7.00$, $df = 25$, $P < 0.0001$; Fig. 6).

Influence of diet on relative rates of shell and tissue growth

In none of the experiments did phytoplankton species affect the relationship between shell and tissue weight, either for larvae or for juveniles, even when shell growth rates differed significantly and substantially within an experiment (Fig. 7). Results were quite different for larvae that had been reared at different concentrations of T-ISO, however. Although larvae reared at the two T-ISO concentrations had similar mean shell weights shortly before metamorphosis ($t = 0.708$, $df = 4$, $P = 0.52$), mean individual tissue weight for larvae reared at the low T-ISO concentration was only 38.6 % that of larvae that had been reared at the high T-ISO concentration ($t = 11.15$, $df = 4$, $P = 0.0004$; Fig. 8), indicating a slower rate of tissue growth relative to shell growth.




Fig. 7 Effect of diet on the relative growth of shell and tissue in *C. fornicata*. The number of replicates is shown within each bar. Each replicate included 4–10 larvae or 1 juvenile. In Experiment 2, larvae had been reared on a diet of DUN for either 3 d, 6 d, or 9 d before

being transferred to a diet of T-ISO prior to metamorphosis; in Experiment 3, larvae were reared on DUN for either 5 or 10 d before being transferred to a diet of T-ISO prior to metamorphosis. See Table 1 for sizes of larvae and juveniles used for these determinations




Fig. 8 Effect of food concentration on relative growth of shell and tissue in larval *C. fornicata*. Larvae were reared on T-ISO until metamorphosis at either 18×10^4 or 1×10^4 cells ml^{-1} , and then raised as juveniles at 18×10^4 cells ml^{-1} . Each mean (+SD) is the average of 3 replicates of 11–15 larvae each. *Significant difference between means

Discussion

As expected from previous research (Klinzing and Pechenik 2000), larvae of *C. fornicata* grew most rapidly on a diet of T-ISO in all experiments. Similarly, larvae of the marine mudsnail *Nassarius obsoletus* (= *Ilyanassa obsoleta*) also grew more quickly when reared on *Isochrysis galbana* (clone ISO) than on *Dunaliella tertiolecta* (DUN; Pechenik and Fisher 1979). Food-specific growth rates could be caused by differences in ingestion rates, digestive rates, or assimilation efficiency, or by subtle differences in nutritional chemistry (Pechenik and Fisher 1979; reviewed by Pechenik 1987). Based on studies with mudsnail larvae (Pechenik and Fisher 1979), in which growth rates, ingestion rates, and assimilation efficiencies were measured for larvae reared on 3 different diets, differences in nutritional content seem the most likely sources of the diet-related differences in mean growth rates seen in the present study. T-ISO is known to be especially rich in certain essential fatty acids (Volkman et al. 1989; Yoshioka et al. 2012), which play a critical role in determining the nutritional value of phytoplankton (Rossoll et al. 2012). Phytoplankton species also vary substantially in vitamin content (Brown et al. 1997). The dietary requirements of *C. fornicata* during larval development have yet to be determined.

There is no reason to suspect that either of the 2 poorer microalgal diets used in this study are toxic to *C. fornicata*. Larval mortality was generally low (<7 %) on all diets, and larval growth rates were linear over time (e.g. Fig. 2). Although mortality was high for larvae reared on DUN in

Experiment 1, it was low in subsequent experiments, arguing against toxicity. Moreover, the highest mean juvenile growth rates recorded in Experiment 2, in which the T-ISO culture had become contaminated with DUN towards the end of the larval rearing period, were comparable to those recorded in Experiments 3 and 5, in which the T-ISO cultures remained pure.

Mean larval growth rates on each diet varied considerably among experiments in our study. Such differences cannot be due to differential larval mortality, as larval mortality was generally low, as already noted, but instead are likely to have a substantial genetic basis; when larvae from 90 females of *C. fornicata* were reared on *Isochrysis galbana*, mean larval growth rates varied from $35 \mu\text{m d}^{-1}$ to over $90 \mu\text{m d}^{-1}$, with a highly significant family effect on larval growth (Hilbish et al. 1999). The remarkably poor growth and survival of larvae on a diet of DUN in our Experiment 1 suggests that larvae from that hatch had unusually high requirements for some particular micronutrient that was either missing from that diet or present at very low concentrations. What those micronutrients are remains to be determined. We noted food in the stomachs of all larvae feeding on all diets, so differences in growth rates were not due to difficulties in capturing phytoplankton cells. There was negligible mortality for larvae being reared on the other 2 phytoplankton species in that experiment, indicating that the larvae were generally healthy, and larval mortality was negligible when we reared larvae on DUN in our other experiments. On the other hand, the variation in mean larval growth rates seen among our experiments on a given diet could also reflect maternal effects (reviewed by Marshall et al. 2008) or differences in algal chemistry among experiments; the fatty acid composition of particular phytoplankton species is known to vary with culture conditions and with the timing of algal harvesting (Chuecas and Riley 1969; Brown et al. 1997). However, we drew phytoplankton from several cultures of each species over the course of each experiment, and cell concentrations at harvesting only varied between ~ 1 and 3×10^6 cells ml^{-1} . Larvae grew at constant rates within experiments (e.g., Fig. 2), suggesting that algal chemistry was stable.

Latent effects of larval diet on juvenile growth were seen in a number of our experiments, including Experiment 2. Juvenile growth rate data from that experiment are potentially complicated by the contamination of our T-ISO culture with DUN late in larval development, as noted earlier. However, the reduced growth rates of juveniles from larvae that had been reared on DUN in that experiment were not likely due to contamination of the juvenile growth media with DUN. Individuals that had been reared as larvae on MONO were stimulated to metamorphose 2–4 days after larvae in any of the other treatments, due to especially slow larval growth on that diet; those individuals would therefore

have been exposed to even higher DUN concentrations during their 4 days of juvenile growth than juveniles from any of the other treatments, and yet their mean juvenile growth rates were the same as those of juveniles that had been raised as larvae on T-ISO. Moreover, as mentioned earlier, mean juvenile growth rates on the contaminated T-ISO in that experiment were comparable to those reported in Hilbish et al. 1999 and other studies. The data suggest, then, that the reduced juvenile growth rates recorded for the DUN treatments in Experiment 2 reflect latent effects of algal diet during larval development, as in Experiment 3, rather than contamination of the T-ISO culture with DUN during the period of juvenile growth.

Perhaps the most interesting finding from this study is that dietary differences during larval development significantly influenced juvenile growth rates in some experiments, but not in others. In Experiment 3, for example, feeding larvae on a diet of MONO through metamorphosis produced latent effects on juvenile growth (Fig. 4B), whereas that was not the case in Experiments 1, 2, or 4, even though larvae grew substantially and significantly more slowly on a diet of MONO than on T-ISO in all of those experiments. Similarly, rearing larvae on a diet of DUN for 9 days influenced post-metamorphic growth rates in Experiment 2 (Fig. 3B), but had no effect on post-metamorphic growth rates in Experiment 3. In previous studies, several days or more of starvation or salinity stress during larval development also produced latent effects on juvenile growth rates in some experiments but not in others (Pechenik et al. 1996a, b, 2002; Bashevkin and Pechenik 2015). Such variability may reflect maternal effects (Marshall et al. 2008)—recall that adults were collected at different times of year for different experiments—or differences in genetic contributions from different parents (Hilbish et al. 1999).

Overall, our data suggest that the effects of larval diet on juvenile development seen in the present study are somehow due to subtle deficiencies in one or more key nutrients being provided to the larvae, and that the larvae of some parents may have somewhat higher requirements for particular fatty acids or other micronutrients than the larvae of other parents. It is important to remember that in our studies, documented latent effects on juvenile growth could not have been caused by genotype-specific differential mortality, since juvenile mortality was generally quite low (<7 %), as in our previous studies with this species (e.g., Klinzing and Pechenik 2000; Pechenik et al. 1996c, 2002).

Note that the larvae in our experiments were triggered to metamorphose soon after they became competent to do so. However, competent larvae of this species can delay their metamorphosis for a considerable time (Pechenik 1980; Pechenik and Eyster 1989), the potential delay period varying inversely with rearing temperature (Zimmerman and

Pechenik 1991). Thus larvae could experience nutritional stress for an even longer period than that which was examined in this study, the consequences of which could be explored in future work.

Despite the diet-related differences in larval growth rates that were seen in Experiments 1–4, and at least sometimes in subsequent juvenile growth rates as well, the proportional contribution of tissue to total dry weight was not altered in any of those experiments, indicating comparable shifts in rates of tissue and shell growth. In contrast, differences in the concentration of T-ISO during larval rearing did alter that relationship—not by altering individual shell weight, but rather by altering individual tissue weight. In studies by Bashevkin and Pechenik (2015), rearing larvae of this same species at different salinities (20 or 30) did not affect relative rates of shell and tissue growth at 25 °C, despite dramatic differences in mean shell growth rates, but did severely impact relative rates of shell and tissue growth at 20 °C. Note that in our study, the effect of reduced food concentration was a relative reduction in tissue growth, whereas reduced salinity at 20 °C caused a relative reduction in shell growth (Bashevkin and Pechenik 2015). We are not aware of any research that examines how the growth of shell and tissue is coordinated during molluscan development, but the topic seems worth exploring.

This study shows that feeding larvae on different microalgal species can indeed negatively impact growth after metamorphosis. Rearing larvae on the best diet (T-ISO) at the low concentration of 1×10^4 cells ml^{-1} throughout larval development also resulted in pronounced reductions in mean juvenile growth rates following metamorphosis, lasting for at least 6 days (Fig. 6). Such latent effects have now been demonstrated for a variety of animals from many phyla in response to a variety of stresses experienced during larval development, including thermal and salinity stress (Pechenik et al. 2001; Thiyagarajan et al. 2007; Wu et al. 2012; Hartmann et al. 2013; Hopkins et al. 2014; Jonsson and Jonsson 2014; Bashevkin and Pechenik 2015), heavy metal exposure (Ng and Keough 2003), ocean acidification (Hettinger et al. 2012, 2013), hypoxia (Li and Chiu 2013), nutritional stress (Pechenik et al. 1996b, 2002, present study; Merilä and Svensson 1997; Phillips 2002; Gardner et al. 2009; Van Allen and Rudolf 2013; Jonsson and Jonsson 2014), delayed metamorphosis (Pechenik et al. 1993; Gebauer et al. 1999; Marshall et al. 2003; Wendt 1998; De Block and Stoks 2005; Thiyagarajan et al. 2007; Graham et al. 2013), and the presence of predators (Relyea and Hoverman 2003; Nicieza et al. 2006; Tejedo et al. 2010; reviewed by Pechenik 2006; Jonsson and Jonsson 2014). Some of these stresses have produced latent effects in some species but not in others. For example, exposing larvae to reduced salinity for 24–48 h caused latent effects for barnacles (Thiyagarajan et al. 2007) and for the polychaete *Capitella teleta* (Pechenik and Cerulli 1991),

but not for *Crepidula fornicata*, even when larval growth rates were severely depressed (Diederich et al. 2011; but see Bashevkin and Pechenik 2015 regarding the impact of longer exposure to salinity stress). Similarly, delayed metamorphosis produced latent effects for the polychaete *Capitella teleta*, the bryozoan *Bugula neritina* (Wendt, 1998), the barnacle *Balanus amphitrite* (Thiyagarajan et al. 2007), and the colonial seasquirt *Diplosoma listerianum* (Marshall et al. 2003), but not for *Crepidula fornicata* (Pechenik and Eyster 1989) or for the solitary ascidian *Styela plicata* (Thiyagarajan and Qian 2003). In addition to the latent effects that have been so widely documented as resulting from some stresses experienced during larval development, it now seems that even stresses experienced by sperm prior to fertilization can influence offspring performance (Ritchie and Marshall 2013). The mechanisms accounting for such latent effects, and for the higher or lower vulnerability of some species—and apparently of the offspring of some parents within a species—to certain stresses, are not yet clear (reviewed by Pechenik 2006). For juvenile *Crepidula fornicata* (Pechenik et al. 2002) and *C. onyx* (Chiu et al. 2007, 2008), reduced juvenile growth seems to be at least partly caused by reduced individual filtration rates, but patterns of transcription, translation, and DNA methylation may also be affected. Molecular studies should help us to determine why some stresses experienced during larval development produce latent effects while others do not, and why the responses differ among species and among the offspring of different parents.

The results of these studies also raise another question: how much of the variation that we see in the natural world is caused by direct genetic differences among individuals and how much reflects the impact of experiences in early development? Our results also add an intriguing note of complexity to attempts at predicting future consequences of environmental change on marine community structure and species interactions, as worldwide phytoplankton concentrations decline (Montes-Hugo et al. 2009; Boyce et al. 2010) and as phytoplankton species composition and nutritional characteristics change (Burkhardt and Riebesell 1997; Rossoll et al. 2012).

Acknowledgments We thank Kelly Boisvert and Melissa MacEwen for their careful assistance in data collection during parts of this study, and two reviewers for their helpful comments and suggestions.

References

- Anger K (1995) Starvation resistance in larvae of a semi-terrestrial crab, *Sesarma curacaoense* (Decapoda: Grapsidae). J Exp Mar Biol Ecol 187:161–174. doi:10.1016/0022-0981(94)00178-G
- Basch LV (1996) Effects of algal and larval densities on development and survival of asteroid larvae. Mar Biol 126:693–701. doi:10.1007/BF00351336
- Bashevkin SM, Pechenik JA (2015) The interactive influence of temperature and salinity on larval and juvenile growth in the gastropod *Crepidula fornicata*. J Exp Mar Biol Ecol 470:78–91. doi:10.1016/j.jembe.2015.05.004
- Blanchard M (1997) Spread of the slipper limpet *Crepidula fornicata* (L. 1758) in Europe. Current state and consequences. Sci Mar 61(Suppl 2):109–118
- Bohn K, Richardson C, Jenkins S (2012) The invasive gastropod *Crepidula fornicata*: reproduction and recruitment in the intertidal at its northernmost range in Wales, UK, and implications for its secondary spread. Mar Biol 159:2091–2103. doi:10.1007/s00227-012-1997-3
- Boyce DG, Lewis MR, Worm B (2010) Global phytoplankton decline over the past century. Nature 466:591–596. doi:10.1038/nature09268
- Brown MR, Jeffrey SW, Volkman JK, Dunstan GA (1997) Nutritional properties of microalgae for mariculture. Aquaculture 151:315–331. doi:10.1016/S0044-8486(96)01501-3
- Burkhardt S, Riebesell U (1997) CO₂ availability affects elemental composition (C:N:P) of the marine diatom *Skeletonema costatum*. Mar Ecol Prog Ser 155:67–76. doi:10.3354/meps155067
- Byrne M (2011) Impact of ocean warming and ocean acidification on marine invertebrate life history stages: vulnerabilities and potential for persistence in a changing ocean. Oceanogr Mar Biol Annu Rev 49:1–42. doi:10.1201/b11009-2
- Chiu JMY, Ng TYT, Wang WX, Thiyagarajan V, Qian PY (2007) Latent effects of larval food limitation on filtration rate, carbon assimilation and growth in juvenile gastropod *Crepidula onyx*. Mar Ecol Prog Ser 343:173–182. doi:10.3354/meps06928
- Chiu JMY, Wang H, Thiyagarajan V, Qian PY (2008) Differential timing of larval starvation effects on filtration rate and growth in juvenile *Crepidula onyx*. Mar Biol 154:91–98. doi:10.1007/s00227-007-0902-y
- Chuecas L, Riley JP (1969) Component fatty acids of the total lipids of some marine phytoplankton. J Mar Biol Assoc UK 49:97–116. doi:10.1017/S0025315400046439
- Cowen RK, Sponaugle S (2009) Larval dispersal and marine population connectivity. Annu Rev Mar Sci 1:443–466. doi:10.1146/annurev.marine.010908.163757
- Crisp DJ (1974) Factors influencing the settlement of marine invertebrate larvae. In: Grant PT, Mackie AM (eds) Chemoreception in marine organisms. Academic Press, New York, pp 177–265
- De Block M, Stoks R (2005) Fitness effects from egg to reproduction: bridging the life history transition. Ecology 86:185–197
- Diederich CM, Jarrett JN, Chaparro OR, Segura CJ, Arellano SM, Pechenik JP (2011) Low salinity stress experienced by larvae does not affect post-metamorphic growth or survival in three calyptraeid gastropods. J Exp Mar Biol Ecol 397:94–105. doi:10.1016/j.jembe.2010.11.019
- Eyster LS, Pechenik JA (1988) Comparison of growth, respiration, and feeding of juvenile *Crepidula fornicata* (L.) following natural or KCl-triggered metamorphosis. J Exp Mar Biol Ecol 118:269–279. doi:10.1016/0022-0981(88)90078-0
- Gardner DS, Ozanne SE, Sinclair KD (2009) Effect of the early-life nutritional environment on fecundity and fertility of mammals. Philos Trans R Soc B 364:3419–3427. doi:10.1098/rstb.2009.0121
- Gebauer P, Paschke K, Anger K (1999) Costs of delayed metamorphosis: reduced growth and survival in early juveniles of an estuarine grapsid crab, *Chasmagnathus granulata*. J Exp Mar Biol Ecol 238:271–281. doi:10.1016/S0022-0981(98)00219-6
- Gosselin LA, Qian PY (1997) Juvenile mortality in benthic marine invertebrates. Mar Ecol Prog Ser 146:265–282. doi:10.3354/meps146265
- Graham EM, Baird AH, Willis BL, Connolly SR (2013) Effects of delayed settlement on post-settlement growth and survival of

- scleractinian coral larvae. *Oecologia* 173:431–438. doi:10.1007/s00442-013-2635-6
- Guillard RRL, Ryther JH (1962) Studies of marine planktonic diatoms. I. *Cyclotella nana* Hustedt, and *Detonula confervacea* (Cleve) Gran. *Can J Microbiol* 8:229–239
- Hansen B (1993) Aspects of feeding, growth and stage development by trochophora larvae of the boreal polychaete *Mediomastus fragile* (Rasmussen)(Capitellidae). *J Exp Mar Biol Ecol* 166:273–288. doi:10.1016/0022-0981(93)90224-C
- Hartmann AC, Marhaver KL, Chamberland VF, Sandin SA, Vermeij MJA (2013) Large birth size does not reduce negative latent effects of harsh environments across life stages in two coral species. *Ecology* 94:1966–1976. doi:10.1890/13-0161.1
- Hayes GG, Richardson AJ, Robinson C (2005) Climate change and marine plankton. *Trends Ecol Evol* 20:337–344. doi:10.1016/j.tree.2005.03.004
- Hettinger A, Sanford E, Hill TM et al (2012) Persistent carry-over effects of planktonic exposure to ocean acidification in the Olympia oyster. *Ecology* 93:2758–2768. doi:10.1890/12-0567.1
- Hettinger A, Sanford E, Hill TM, Lenz EA, Russell AD, Gaylord B (2013) Larval carry-over effects from ocean acidification persist in the natural environment. *Glob Change Biol* 19:3317–3326. doi:10.1111/gcb.12307
- Hilbish TJ, Sasada K, Eyster LS, Pechenik JA (1999) Relationship between rates of swimming and growth in veliger larvae: genetic variance and covariance. *J Exp Mar Biol Ecol* 239:183–193. doi:10.1016/S0022-0981(99)00009-X
- Hinga KR (2002) Effects of pH on coastal marine phytoplankton. *Mar Ecol Prog Ser* 238:281–300. doi:10.3354/meps238281
- His E, Seaman MNL (1992) Effects of temporary starvation on the survival, and on subsequent feeding and growth, of oyster (*Crassostrea gigas*) larvae. *Mar Biol* 114:277–279. doi:10.1007/BF00349530
- Hoogstraten A, Timmermans KR (2012) Morphological and physiological effects in *Proboscia alata* (Bacillariophyceae) grown under different light and CO₂ conditions of the modern Southern Ocean. *J Phycol* 48:559–568. doi:10.1111/j.1529-8817.2012.01148.x
- Hopkins GR, Brodie ED Jr, French SS (2014) Developmental and evolutionary history affect survival in stressful environments. *PLoS ONE* 9:e95174. doi:10.1371/journal.pone.0095174
- Hunt HL, Scheibling RE (1997) Role of early post-settlement mortality in recruitment of benthic marine invertebrates. *Mar Ecol Prog Ser* 155:269–301. doi:10.3354/meps155269
- Jennings LB, Hunt HL (2011) Small macrobenthic invertebrates affect the mortality and growth of early post-settlement sea urchins and sea stars in subtidal cobble habitat. *Mar Ecol Prog Ser* 431:173–182. doi:10.3354/meps09131
- Jonsson B, Jonsson N (2014) Early environment influences later performance in fishes. *J Fish Biol* 85:151–188. doi:10.1111/jfb.12432
- Kim J-M, Lee K, Shin K, Kang J-H, Lee H-W, Kim M, Jang P-G, Jang M-C (2006) The effect of seawater CO₂ concentration on growth of a natural phytoplankton assemblage in a controlled mesocosm experiment. *Limnol Oceanogr* 51:1629–1636. doi:10.4319/lo.2006.51.4.1629
- Klinzing MS, Pechenik JA (2000) Evaluating whether velar-lobe size indicates food limitation among larvae of the marine gastropod *Crepidula fornicata*. *J Exp Mar Biol Ecol* 252:255–279. doi:10.1016/S0022-0981(00)00245-8
- Leu E, Daase M, Schulz KG, Stühr A, Riebesell U (2013) Effect of ocean acidification on the fatty acid composition of a natural plankton community. *Biogeosciences* 10:1143–1153. doi:10.5194/bg-10-1143-2013
- Li A, Chiu JMY (2013) Latent effects of hypoxia on the gastropod *Crepidula onyx*. *Mar Ecol Prog Ser* 480:145–154. doi:10.3354/meps10213
- Lucas JS, Costlow JD Jr (1979) Effects of various temperature cycles on the larval development of the gastropod mollusc *Crepidula fornicata*. *Mar Biol* 51:111–117. doi:10.1007/BF00555190
- Marshall DJ, Pechenik JA, Keough MJ (2003) Larval activity levels and delayed metamorphosis affect post-larval performance in the colonial ascidian *Diplosoma listerianum*. *Mar Ecol Prog Ser* 246:153–162. doi:10.3354/meps246153
- Marshall DJ, Allen RM, Crean AJ (2008) The ecological and evolutionary importance of maternal effects in the sea. *Oceanogr Mar Biol Ann Rev* 46:203–250
- McEdward LR, Qian P-Y (2001) Effects of the duration and timing of starvation during larval life on the metamorphosis and initial size of the polychaete *Hydroides elegans* (Haswell). *J Exp Mar Biol Ecol* 261:185–197. doi:10.1016/S0022-0981(01)00272-6
- Merilä J, Svensson E (1997) Are fat reserves in migratory birds affected by condition in early life? *J Avian Biol* 28:279–286. doi:10.2307/3676940
- Montes-Hugo M, Doney SC, Ducklow HW, Fraser W, Martinson D, Stammerjohn SE, Schofield O (2009) Recent changes in phytoplankton communities associated with rapid regional climate change along the Western Antarctic Peninsula. *Science* 323:1470–1473. doi:10.1126/science.1164533
- Morgan SG (1995) Life and death in the plankton: larval mortality and adaptation. In: McEdward L (ed) *Ecology of marine invertebrate larvae*. CRC Press, New York, pp 279–321
- Ng TYT, Keough MJ (2003) Delayed effects of larval exposure to Cu in the bryozoan *Watersipora subtorquata*. *Mar Ecol Prog Ser* 257:77–85. doi:10.3354/meps257077
- Nicieza AG, Álvarez D, Atienza EMS (2006) Delayed effects of larval predation risk and food quality on anuran juvenile performance. *J Evol Biol* 19:1092–1103. doi:10.1111/j.1420-9101.2006.01100.x
- O'Connor MI, Bruno JF, Gaines SD, Halpern BS, Lester SE, Kinlan BP, Weiss JM (2007) Temperature control of larval dispersal and the implications for marine ecology, evolution, and conservation. *Proc Natl Acad Sci USA* 104:1266–1271
- Paine RT (1964) Ash and calorie determinations of sponge and opisthobranch tissues. *Ecology* 45:384–387. doi:10.2307/1933856
- Pechenik JA (1980) Relationship between rate of development and duration of larval life in larvae of the marine prosobranch gastropod *Crepidula fornicata*. *J Exp Mar Biol Ecol* 74:241–257
- Pechenik JA (1987) Environmental influences on larval survival and growth. In: Giese AC, Pearse JS (eds) *Reproduction of Marine Invertebrates*, vol 9. Blackwell, New York, pp 551–608
- Pechenik JA (1999) On the advantages and disadvantages of larval stages in benthic marine invertebrate life cycles. *Mar Ecol Prog Ser* 177:269–297. doi:10.3354/meps177269
- Pechenik JA (2006) Larval experience and latent effects—metamorphosis is not a new beginning. *Integr Comp Biol* 46:323–333. doi:10.1093/icb/icj028
- Pechenik JA, Cerulli T (1991) Influence of delayed metamorphosis on survival, growth, and reproduction of the marine polychaete *Capitella* sp. I. *J Exp Mar Biol Ecol* 151:17–27. doi:10.1016/0022-0981(91)90012-L
- Pechenik JA, Eyster LS (1989) Influence of delayed metamorphosis on the growth and metabolism of young *Crepidula fornicata* (Gastropoda) juveniles. *Biol Bull* 176:14–24. doi:10.2307/1541884
- Pechenik JA, Fisher NS (1979) Ingestion and assimilation of three phytoplankton species related to larval growth rates of the mud snail, *Nassarius obsoletus*. *J Exp Mar Biol Ecol* 38:57–80

- Pechenik JA, Gee CC (1993) Onset of metamorphic competence in larvae of the gastropod *Crepidula fornicata*, judged by a natural and an artificial cue. *J Exp Mar Biol Ecol* 167:59–72
- Pechenik JA, Heyman WD (1987) Using KCl to determine size at competence for larvae of the marine gastropod *Crepidula fornicata* (L.). *J Exp Mar Biol Ecol* 112:27–38. doi:10.1016/S0022-0981(87)80012-6
- Pechenik JA, Rittschof D, Schmidt AR (1993) Influence of delayed metamorphosis on survival and growth of juvenile barnacles *Balanus amphitrite*. *Mar Biol* 115:287–294. doi:10.1007/BF00346346
- Pechenik JA, Estrella S, Hammer K (1996a) Food limitation stimulates metamorphosis and alters postmetamorphic growth rate in the marine prosobranch gastropod *Crepidula fornicata*. *Mar Biol* 127:267–275
- Pechenik JA, Hammer K, Weise C (1996b) The effect of starvation on acquisition of competence and postmetamorphic performance in the marine prosobranch gastropod *Crepidula fornicata* (L.). *J Exp Mar Biol Ecol* 199:137–152. doi:10.1016/0022-0981(96)00010-X
- Pechenik JA, Hilbish TJ, Eyster LS, Marshall D (1996c) Relationship between larval and juvenile growth rates in two marine gastropods, *Crepidula plana* and *C. fornicata*. *Mar Biol* 125:119–127. doi:10.1007/BF00350766
- Pechenik JA, Gleason T, Daniels D, Champlin D (2001) Influence of larval exposure to salinity and cadmium stress on juvenile performance of two marine invertebrates (*Capitella sp. I* and *Crepidula fornicata*). *J Exp Mar Biol Ecol* 264:101–114. doi:10.1016/S0022-0981(01)00313-6
- Pechenik JA, Jarrett J, Rooney J (2002) Relationship between larval nutritional experience, larval growth rates, and juvenile growth rates in the prosobranch gastropod *Crepidula fornicata*. *J Exp Mar Biol Ecol* 280:63–78. doi:10.1016/S0022-0981(02)00367-2
- Pechenik JA, Ambrogio OV, Untersee S (2010) Predation on juveniles of *Crepidula fornicata* by two crustaceans and two gastropods. *J Exp Mar Biol Ecol* 384:91–98. doi:10.1016/j.jembe.2009.12.011
- Phillips NE (2002) Effects of nutrition-mediated larval condition on juvenile performance in a marine mussel. *Ecology* 83:2562–2572
- Pires A, Guilbault TR, Mitten JV, Skiendzielewski JA (2000) Catecholamines in larvae and juveniles of the prosobranch gastropod, *Crepidula fornicata*. *Comp Biochem Physiol C* 127:37–47. doi:10.1016/S0742-8413(00)00128-6
- Relyea RA, Hoverman JT (2003) The impact of larval predators and competitors on the morphology and fitness of juvenile treefrogs. *Oecologia* 134:596–604. doi:10.1007/s00442-002-1161-8
- Ritchie H, Marshall DJ (2013) Fertilisation is not a new beginning: sperm environment affects offspring developmental success. *J Exp Biol* 216:3104–3109. doi:10.1242/jeb.087221
- Rossoll D, Bermúdez R, Hauss H, Schulz KG, Riebesell U, Sommer U, Winder M (2012) Ocean acidification-induced food quality deterioration constrains trophic transfer. *PLoS ONE* 7:e34737. doi:10.1371/journal.pone.0034737
- Rumrill SS (1990) Natural mortality of marine invertebrate larvae. *Ophelia* 32:163–198. doi:10.1080/00785236.1990.10422030
- Tejedo M, Marangoni F, Pertoldi C, Richter-Boix A, Laurila A, Orizaola G, Nicieza AG, Álvarez D, Gomez-Mestre I (2010) Contrasting effects of environmental factors during larval stage on morphological plasticity in post-metamorphic frogs. *Clim Res* 43:31–39. doi:10.3354/cr00878
- Thiyagarajan V, Qian PY (2003) Effect of temperature, salinity and delayed attachment on development of the solitary ascidian *Styela plicata* (Lesueur). *J Exp Mar Biol Ecol* 290:133–146. doi:10.1016/S0022-0981(03)00071-6
- Thiyagarajan V, Pechenik JA, Gosselein LA, Qian PY (2007) Juvenile growth in barnacles: combined effect of delayed metamorphosis and sub-lethal exposure of cyprids to low-salinity stress. *Mar Ecol Prog Ser* 344:173–184. doi:10.3354/meps06931
- Thorson G (1950) Reproductive and larval ecology of marine bottom invertebrates. *Biol Rev* 25:1–45. doi:10.1111/j.1469-185X.1950.tb00585.x
- Tortell PD, DiTullio GR, Sigman DM, Morel FMM (2002) CO₂ effects on taxonomic composition and nutrient utilization in an Equatorial Pacific phytoplankton assemblage. *Mar Ecol Prog Ser* 236:37–43. doi:10.3354/meps236037
- Van Allen BG, Rudolf VHW (2013) Ghosts of habitats past: environmental carry-over effects drive population dynamics in novel habitat. *Am Nat* 181:596–608. doi:10.1086/670127
- Vaughn D, Allen JD (2010) The peril of the plankton. *Integr Comp Biol* 50:552–570. doi:10.1093/icb/icq037
- Volkman JK, Jeffrey SW, Nichols PD, Rogers GI, Garland CD (1989) Fatty acid and lipid composition of 10 species of microalgae used in mariculture. *J Exp Mar Biol Ecol* 128:219–240. doi:10.1016/0022-0981(89)90029-4
- Wendt DE (1998) Effect of larval swimming duration on growth and reproduction of *Bugula neritina* (Bryozoa) under field conditions. *Biol Bull* 195:126–135. doi:10.2307/1542820
- Wu C-S, Gomez-Mestre I, Kam Y-C (2012) Irreversibility of a bad start: early exposure to osmotic stress limits growth and adaptive developmental plasticity. *Oecologia* 169:15–22. doi:10.1007/s00442-011-2170-2
- Wynne-Edwards C, King R, Davidson A, Wright S, Nichols PD, Wotherspoon S, Kawaguchi S, Virtue P (2014) Species-specific variations in the nutritional quality of southern ocean phytoplankton in response to elevated pCO₂. *Water* 6:1840–1859
- Yoshioka M, Yago T, Yoshie-Stark Y, Arakawa H (2012) Effect of high frequency of intermittent light on the growth and fatty acid profile of *Isochrysis galbana*. *Aquaculture* 338–341:111–117. doi:10.1016/j.aquaculture.2012.01.005
- Zimmerman KM, Pechenik JA (1991) How do temperature and salinity affect relative rates of growth, morphological differentiation, and time to metamorphic competence in larvae of the marine gastropod *Crepidula plana*? *Biol Bull* 180:372–386