

# Vasculature of the hive: heat dissipation in the honey bee (*Apis mellifera*) hive

Rachael E. Bonoan · Rhyan R. Goldman ·  
Peter Y. Wong · Philip T. Starks

Received: 24 February 2014 / Revised: 1 April 2014 / Accepted: 3 April 2014 / Published online: 24 April 2014  
© Springer-Verlag Berlin Heidelberg 2014

**Abstract** Eusocial insects are distinguished by their elaborate cooperative behavior and are sometimes defined as superorganisms. As a nest-bound superorganism, individuals work together to maintain favorable nest conditions. Residing in temperate environments, honey bees (*Apis mellifera*) work especially hard to maintain brood comb temperature between 32 and 36 °C. Heat shielding is a social homeostatic mechanism employed to combat local heat stress. Workers press the ventral side of their bodies against heated surfaces, absorb heat, and thus protect developing brood. While the absorption of heat has been characterized, the dissipation of absorbed heat has not. Our study characterized both how effectively worker bees absorb heat during heat shielding, and where worker bees dissipate absorbed heat. Hives were experimentally heated for 15 min during which internal temperatures and heat shielder counts were taken. Once the heat source was removed, hives were photographed with a thermal imaging camera for 15 min. Thermal images allowed for spatial tracking of heat flow as cooling occurred. Data indicate that honey bee workers collectively minimize heat gain during heating and accelerate heat loss during cooling. Thermal images show that heated areas temporarily increase in size in all directions and then rapidly decrease to safe levels (<37 °C). As such, heat shielding is reminiscent of bioheat removal via the cardiovascular system of mammals.

**Keywords** Colonies · Group behavior · Heat transfer · Temperature dynamics

## Introduction

Social insects are the most abundant of land-dwelling arthropods and are found, among other habitats, in almost all forests around the world (Hölldobler and Wilson 2009). Their success and diversity is attributed to the division of labor and cooperative behavior exhibited within these social groups. Eusocial insects (i.e., ants, termites, and colonial wasps and bees) exhibit such extreme levels of cooperation that the colony is sometimes classified as a superorganism. A superorganism (as first defined in reference to ants) is a group of individuals that collectively share the characteristics of an organism (Wheeler 1910; Tautz 2008; Hölldobler and Wilson 2009). In this regard, each individual in the colony is analogous to a single cell and each caste is analogous to an organ system (Wheeler 1910; Anderson and McShea 2001; Tautz 2008; Hölldobler and Wilson 2009).

The nest serves as the skin and skeleton of this superorganism. The nest provides a microhabitat that allows for social life to happen—it is where food is stored, brood are raised, and colony members interact. As such, the nest must be appropriately protected and maintained—we call behaviors designed to accomplish this social homeostatic mechanisms (Wilson 1971). As the nest is both built and maintained via a collective effort, research on nest architecture and social homeostasis is integral to understanding the evolution of social behavior (Hansell 1996).

Temperature maintenance is a social homeostatic mechanism that lends itself well to experimentation. Particularly important to brood development, temperature is consistently maintained in a variety of social insect nests, despite their structural diversity. The first step in creating a buffer between

Communicated by: Sven Thatje

Rachael E. Bonoan and Rhyan R. Goldman contributed equally to this work.

R. E. Bonoan · R. R. Goldman · P. T. Starks (✉)  
Department of Biology, Tufts University, 163 Packard Ave, Medford,  
MA, USA  
e-mail: philip.starks@tufts.edu

P. Y. Wong  
Department of Mechanical Engineering, Tufts University, 200  
College Ave, Medford, MA, USA

ambient and nest temperatures is selecting a nest location (Jones and Oldroyd 2006). As such, location is very much a part of the nest. Honey bees nest in cavities that provide insulation (Heinrich 1979; Jones and Oldroyd 2006), while termites and many ant species build intricate underground nests equipped with ventilation (Wheeler 1910; Korb 2003; Jones and Oldroyd 2006; Hölldobler and Wilson 2009). Social wasps construct enveloped and unenveloped nests in diverse locations including cavities found both above and below ground (Jeanne and Morgan 1992; Jones and Oldroyd 2006).

When ambient temperatures are not favorable for brood development, both ants and termites actively move brood to more protected areas (Wheeler 1910; Wilson 1971; Korb 2003; Hölldobler and Wilson 2009). In contrast, honey bees and social wasps cannot physically move their brood and must actively regulate temperature. For example, both honey bees and social wasps use evaporative cooling when ambient temperatures are too high for proper brood development (Wilson 1971; Prange 1996). Despite these similarities, honey bees stand alone when it comes to controlling even the slightest temperature fluctuations within their nest.

As part of a colony, adult honey bee workers can withstand hive temperatures up to 50 °C (Coelho 1991), but brood must remain between 32 and 36 °C with a specific preferred temperature of  $34.5 \pm 1.5$  °C (Kronenberg and Heller 1982; Jones et al. 2005; Tautz 2008). This temperature range is necessary for proper larval as well as pupal development (Kronenberg and Heller 1982; Winston 1987). Temperatures higher than 36 °C can increase brood mortality, delay development time, and cause malformations of the wings, stinger, and proboscis (Fukuda and Sakagami 1968; Winston 1987; Groh et al. 2004). Conversely, temperatures lower than 32 °C can result in immune compromise and a decrease in foraging performance as an adult (Winston 1987; Tautz et al. 2003; Groh et al. 2004; Jones et al. 2005). Temperature is not as strictly maintained in all areas of the hive; stable temperatures are not as necessary for resources such as pollen and honey (Fahrenholz et al. 1989). Even within the brood comb, temperature maintenance varies; pupae are more sensitive to variable temperatures than are larvae (Jones et al. 2005).

To maintain brood comb temperature range during brood development, honey bees use a variety of thermoregulatory behaviors. For example, to increase temperature, workers create heat by isometrically contracting thoracic muscles, similar to shivering in mammals (Heinrich 1980, 1985; Bujok et al. 2002; Kleinhenz et al. 2003). To increase the overall temperature of the brood comb, multiple workers contract their muscles, simultaneously heating many larvae/pupae (Heinrich 1980, 1985). To increase the temperature locally, a single worker can enter an empty cell and warm the adjacent brood (Bujok et al. 2002; Kleinhenz et al. 2003). In order to survive long periods of cold temperature, workers exhibit clustering to maintain heat within the hive. During

clustering, workers huddle together to retain heat actively produced by the workers at the center of the cluster (Simpson 1961; Kronenberg and Heller 1982; Stabentheiner et al. 2003).

During hot conditions, workers are able to decrease temperatures. To decrease the temperature on a large scale, workers fan the hive with their wings and may simultaneously spread water to induce evaporative cooling (Heinrich 1979, 1980, 1985; Prange 1996). To decrease temperature on a fine scale, workers use a behavior called heat shielding. To achieve heat shielding, young workers orient themselves between a heat source and brood comb, creating a physical barrier where they passively absorb heat (Starks and Gilley 1999; Siegel et al. 2005; Starks et al. 2005). Most workers heat shield by placing their ventral side directly against a heated surface (Starks and Gilley 1999). In conjunction, other workers have been observed orienting their ventral surface against potentially affected brood comb (Siegel et al. 2005). Research has shown that heat shielding is a context-dependent response; changes in intensity of heat, placement of heat, and density of brood all influence the number of workers that engage in the behavior (Starks and Gilley 1999; Siegel et al. 2005; Starks et al. 2005).

Once workers have absorbed heat, it must be dissipated away from the brood. Studies have shown that foragers can cool their bodies by flying to simulate wind (Heinrich 1979, 1980; Fahrenholz et al. 1989) and by regurgitating nectar, allowing it to absorb excess heat (Prange 1996). While this provides a mechanism for heat dissipation, it does not provide information on where workers dissipate absorbed heat within a hive. To investigate this particular aspect of social homeostasis, we created localized heat stress in experimental honey bee hives and used thermal imaging to visualize movement following experimental heat stress. At least two types of heat movement are possible: (1) pattern-free dissipation and (2) pattern-rich dissipation. Pattern-free dissipation would be characterized by no trend in the direction in which heated workers move within the hive. Pattern-rich dissipation would show some directed trend in movement of heated workers. This movement might be out of the hive or to a less regulated part of the hive (i.e., where there is stored honey or pollen).

## Materials and methods

### Subjects

In June 2013, seven two-frame Plexiglas (6 mm thick) observation hives (53 cm×48 cm×5 cm) with active honey bee colonies were installed at the Tufts University International Social Insect Research Facility in Medford, MA. Each hive was re-queened before transportation, and queens were restricted to one frame using a queen excluder. This ensured

that for the duration of the experiment, all brood would be laid and reared in only one of the two frames (see Siegel et al. 2005 and Starks et al. 2005; Fig. 1).

After installation, the facility was kept at a constant temperature of approximately 20 °C. For 1 week, the newly installed hives were fed 100 ml of 1:1 sucrose water each day during orientation and acclimation. During the initial feeding period, each hive was censused for approximate number of bees. The approximate colony sizes were determined to be 1,000–2,500 adult bees (Sammataro and Avitabile 2011). Comb maps of capped and uncapped brood, capped and uncapped honey, pollen, and empty cells were generated to identify similar areas across hives for treatment (i.e., heating). Because workers preferentially shield brood comb (Starks and Gilley 1999; Starks et al. 2005), we identified areas in each experimental hive that had similar amounts of brood. A 12.5 cm×10 cm section at the left center of the lower frame was found to have similar quantities of brood comb in all hives and was subjected to heating during experimental trials (Fig. 1).

#### Collection of temperature data during heating


To minimize heating of the surrounding hive areas, hives were fitted with insulation (Foamular 250 1-in. insulation,  $R=5.0$ ). A 12.5-cm×10-cm rectangle was cut out of the lower left quadrant of each piece of insulation (see above). Since bees do not see red light, the experimental window was covered with red theater gel (Daumer 1956, as cited in von Frisch

1967; Gribakin 1969). This ensured that any observed change in behavior would be due to the presence of heat and not the presence of light. To heat the uninsulated experimental window, a theater lamp with a heat bulb (GE 250 W infrared heat reflector bulb) was pre-heated for 5 min (reaching 155.9–173.0 °C) and then placed 50 cm away from the window. From July 11th, 2013 to July 19th, 2013, each of the seven experimental hives and a control hive were heated for 15 min using the methods described above. Two trials were run for each hive; all trials were done between the hours of 9:00 a.m. and 12:00 p.m. Additionally, all trials were run in the dark in order to mimic the natural nest environment and minimize possible light effects.

Internal temperature data were collected before heating, during heating, and immediately following heating. During heating, the temperature—both under the heated window and under the insulation—was taken every minute using an Omega handheld digital thermometer with type K Teflon insulated thermocouples sensitive to 0.1 °C (see numbered thermocouples in Fig. 1).

A control hive—a hive with all comb characteristics but no honey bees—was heated to approximately normal brood comb temperatures ( $31.1\pm 4.7$  °C) using two small electric heating pads (ZooMed Repti-therm 4 W heat pad). Coupled with the presence of the typical comb characteristics (brood, pollen and honey), the heating pads simulated a hive environment as if there were active bees in the hive. Once the control hive was stable at approximately normal brood comb

**Fig. 1** Diagram of the experimental setup and photo of the interior hive structure. Each hive contained two frames separated by a queen excluder. The characteristic pattern of hive structure is made clear by the dashed lines. After surveying all seven experimental hives, the area selected for heating was to the left center of the brood area. This particular area was selected as it had similar brood densities across all hives. For the purposes of temperature collection, one thermocouple (1) was situated on the Plexiglas that was to be covered with the insulation. The second thermocouple (2) was situated in the brood comb of the heated window so that internal temperatures before, during, and immediately following heating could be recorded



temperatures, the hive was subjected to the heating protocol outlined above.

#### Observation of heat shielders

The number of heat shielders—as evidenced by individuals with the ventral side of their bodies placed against the heated Plexiglas—was counted before and immediately after each heating period (see methods in Starks and Gilley 1999; Siegel et al. 2005; Starks et al. 2005).

#### Collection of thermal images during cooling

Immediately after counting heat shielders, the insulation was removed and a thermal imaging camera (Fluke Thermal Imager model Ti32, emissivity=0.95) was used to take infrared images of the hives every 30 s for 15 min. In addition to general heat maps, the camera provided data on temperature extremes on the external Plexiglas surface of the observation hive. This procedure was repeated twice for each hive. Each hive was allowed a full day to recover between trials. Hives were never allowed to heat to an internal temperature above 41 °C and were monitored for detrimental effects (i.e., increased mortality of workers and brood, abnormally slow workers) within the heated window—none were observed.

#### Statistical analysis


Temperature gained during heating was analyzed using a  $2 \times 2$  mixed effect ANOVA (between=control vs. experimental, within=under window vs. under insulation); assumptions of equal variance and normal distribution were met. The change in heat shielder number before ( $n=14$ ) and after ( $n=14$ ) heating was analyzed via a paired  $t$  test. A Welch two-

sample  $t$  test was used to compare control ( $n=2$ ) and experimental ( $n=14$ ) hives for the average temperature post-heating. For each image generated during cooling, the area of heated regions at or above 37 °C was determined (see the red and white areas in Fig. 2). The mean area measured in control ( $n=62$ ) and experimental ( $n=434$ ) hives at each time point was compared using a Welch two-sample  $t$  test. Regression analysis was run for both control ( $n=2$ ) and experimental ( $n=14$ ) hives to determine the rate in temperature decrease post-heating (time period=0–9 min; on average, experimental hives had cooled to 37 °C in 9 min). Statistics were performed using R Version 3.0.2 (R Core Team 2013).

## Results


#### Heating period

During the 15-min heating period, overall temperature gain was significantly greater in control relative to experimental hives ( $F_1=93.43$ ,  $p<0.001$ ; Fig. 3). The average internal temperature gain under the insulation was significantly lower than the average internal temperature gain under the heated window ( $F_1=117.614$ ,  $p<0.001$ ; Fig. 3). This was seen in both control and experimental hives, indicating that the insulation was effective in creating localized heat stress. Consistent with the creation of localized heat stress, workers displayed heat shielding; there were significantly more heat shielders after heating than before ( $t_{13}=2.82$ ,  $p=0.01$ ; Fig. 4). Immediately after heating, the average external temperature of the Plexiglas window was significantly higher for the control hive ( $48.5 \pm 0.8$  °C) than for experimental hives ( $46.5 \pm 0.6$  °C) ( $t_{15}=5.73$ ,  $p<0.001$ ; Fig. 5).



**Fig. 2** Comparison of representative experimental and control infrared images taken pre- and post-heating. These images are in the same orientation as in Figure 1. The *color green* indicates the presences of bees in the experimental hive and the heating pads in the control hive. *Red and white areas* indicate temperatures above 37 °C. In the experimental hive, the *red*

*area* grew significantly larger within 3 min of cooling and disappeared within 9 min. In contrast, the high heat area in the control hive gradually decreased in size and still persisted after 18 min of cooling. Such differences indicate that workers effectively cooled the hive by absorbing the heat moving it into the periphery




**Fig. 3** Mean change in internal temperature for insulated and uninsulated regions of the observation hives. For both control ( $n=2$ ) and experimental hives ( $n=14$ ), the temperature increase was more gradual under the insulation than under the heated window ( $p<0.001$ ). The heat gain in the experimental hive was significantly smaller than the gain in the control hive ( $p<0.001$ ). Since the control hive lacked workers, these data demonstrate that the workers are responsible for regulating temperature changes within the hive. Data were taken from seven experimental hives and one empty control hive. Two trials were done for each hive. Error bars represent one standard error


Cooling period

During the 15-min cooling period, the control hive did not reach temperatures safe for brood development (i.e., below 37 °C; Figs. 2 and 5). By comparison, the mean temperature of the experimental hives reached safe levels within 10 min (Figs. 2 and 5). The mean temperature of the experimental hives post-heating ( $38.7\pm3.4$  °C) was significantly lower than that of the control hives ( $42.5\pm2.5$  °C) ( $t_{98,61}=10.48$ ,  $p<0.001$ ; Fig. 5). On average, the experimental hive cooled to a temperature less than 37 °C almost twice as fast as the control hive (slope  $[0:9 \text{ min}]_{\text{exp}}=-1.87$ , slope  $[0:9 \text{ min}]_{\text{con}}=-1.07$ ; Fig. 5).

When the thermal images of experimental and control hives were compared, differences in the area of regions above 37 °C were observed (Figs. 2 and 6). Immediately after heating, control and experimental hives exhibited a similarly sized heated area (Figs. 2 and 6). However, as cooling




**Fig. 4** Mean number of heat shielders before ( $n=14$ ) and after heating ( $n=14$ ). There were significantly more heat shielders observed after heating than before heating ( $p=0.01$ ) indicating the heat shielding occurred. Data are from seven observation hives, two trials each



**Fig. 5** Mean change in surface temperature of the heated section of brood comb during the 15-min cool down period. Immediately after heating (time=0), the control hive ( $n=2$ ) had a significantly higher mean temperature than the experimental hive ( $n=14$ ;  $p=0.005$ ). After only 10 min of cooling, the experimental hives were able to reach a safe temperature ( $<37$  °C) as highlighted by the shaded gray area. In contrast, the control hive did not reach safe temperatures until after the 15-min cooling period (data not shown). Data were taken from seven experimental hives and one empty control hive with comb. Two trials were done for each hive. Error bars represent one standard error

continued, the area that showed temperatures above 37 °C increased in the experimental hives; no such effect was seen in the control hive ( $t_{372,64}=4.32$ ,  $p<0.001$ ; Fig. 5). By about 3 min post-heating, the average size of the high heat area in the experimental hive had increased dramatically in all directions ( $\Delta \text{area}=6,660\pm2,630 \text{ pixels}^2$ ). Following this spike, the heated area of the experimental hives then decreased rapidly until the hive reached safe levels. By comparison, the high heat regions in the control hive decreased gradually without any increases and had not cooled even after 18 min (Figs. 2 and 6).



**Fig. 6** The average percentage change in the area of the heated region for experimental ( $n=434$ ) and control ( $n=62$ ) hives over time. The heated region was defined as the red area above 37 °C (see Online Resource 1) for each generated heat map. The percent increase in the experimental hives demonstrates a dramatic increase in the high heat area within the first 5 min; this pattern was not observed in the control hive. The high heat area within the experimental hives—but not the control hive—then rapidly decreased until the high heat region disappeared and the hives were cooled to safe levels ( $<37$  °C). Data were from seven observation hives and one control hive with comb, two trials were done for each hive. Error bars represent one standard error

## Discussion

Consistent with previous research, there was a significant increase in the number of heat shielders after heating, indicating that heat shielding occurred (Fig. 4). As such, temperature dynamics within heated regions of the hive were likely influenced by this behavior. In addition, during the 15-min heating period, temperatures remained lower in the heated window of the experimental hives than in the control hive (Fig. 3). Since the control hive did not have bees, differences in temperature were likely caused by the worker bees themselves. The significantly lower temperature gain in the experimental hives highlights the workers' ability to minimize temperature increases during localized heat stress (Fig. 3). Similarly, workers effectively lowered brood comb temperature back to safe levels within 10 min (Fig. 5).

While cooling, the change in area of the high heat regions between the experimental and the control hives differed markedly (Fig. 6). The experimental hives' sudden increase in high heat area—and the lack of an increase in the control hive—implies that the workers were actively moving heat out of the heated region (Fig. 6). Thermal images show that the area increases in *all* directions from the heated point in the hive, showing a radial movement of the workers to the periphery of the hive (Fig. 2). In a natural hive, this movement would drive heat to less regulated areas of the hive, such as honey and pollen stores (Seeley and Morse 1976). Since feral honey bees build their comb in the same vertical fashion that is found in observation hives, our data are representative of what may occur in the field (Winston 1987).

This is the first study to characterize where heat is moved following heat shielding, and how effectively workers dissipate this absorbed heat. The results of this study demonstrate that workers are able to work in concert to inhibit localized temperature gain, and then work rapidly to dissipate what temperature gain is experienced over the developing brood. The initial expansion of the heated area—moving heat from hot to cool areas—is reminiscent of bioheat transfer via the cardiovascular system of mammals. Thus, these data provide additional support for the argument that a honey bee colony can be viewed as a superorganism (Seeley 1989) as well as a concrete example of social homeostasis within a nest.

**Acknowledgments** We thank Anne A. Madden, Kelsey Graham, Tyler Corey, and Georgiana Burruss for providing comments on previous drafts. This research was supported by the Tufts University Biology Department and the Tufts University National Science Foundation Research Experience for Undergraduates Program (DBI 263030).

## References

- Anderson C, McShea DW (2001) Intermediate-level parts in insect societies: adaptive structures that ants build away from the nest. *Insectes soc* 48:291–301
- Bujok B, Kleinhenz M, Fuchs S, Tautz J (2002) Hot spots in the bee hive. *Naturwissenschaften* 89:299–301
- Coelho J (1991) Heat transfer and body temperature in honey bee (Hymenoptera: Apidae) drones and workers. *Environ Entomol* 20:1627–1635
- Daumer K (1956) Metric study of color vision in bees. *J Comp Physiol* 38:413–478
- Fahrenholz L, Lamprecht I, Schrickler B (1989) Thermal investigations of a honey bee colony: thermoregulation of the hive during summer and winter and heat production of members of different bee castes. *J Comp Physiol B* 159:551–560
- Fukuda H, Sakagami SF (1968) Worker brood survival in honeybees. *Res Popul Ecol (Kyoto)* 10:31–39
- Gribakin FG (1969) Cellular basis of colour vision in the honey bee. *Nature* 223:639–641
- Groh C, Tautz J, Rossler W (2004) Synaptic organization in the adult honey bees brain is influenced by brood-temperature control during pupal development. *Proc Natl Acad Sci U S A* 101:4268–4273
- Hansell MH (1996) Wasps make nests: nests make conditions. In: Turillazzi S, West-Eberhard MJ (eds) *Natural history and evolution of paper-wasps*. Oxford University Press, Great Britain, pp 274–289
- Heinrich B (1979) Keeping a cool head: honeybee thermoregulation. *Science* 205:1269–1271
- Heinrich B (1980) Mechanisms of body-temperature regulation in honeybees. *Apis mellifera*: I. Regulation of head temperature. *J Exp Biol* 85:61–72
- Heinrich B (1985) The social physiology of temperature regulation in honeybees. *Fortsch Zool* 31:393–406
- Hölldobler B, Wilson EO (2009) *The superorganism*. W.W. Norton and Company, New York
- Jeanne RL, Morgan RC (1992) The influence of temperature on nest site choice and reproductive strategy in a temperate zone *Polistes* wasp. *Ecol Entomol* 17:135–141
- Jones JC, Helliewell P, Beekman M, Malezka R, Oldroyd BP (2005) The effects of rearing temperature on developmental stability and learning and memory in the honey bee, *Apis mellifera*. *J Comp Physiol A* 191:1121–1129
- Jones JC, Oldroyd BP (2006) Nest thermoregulation in social insects. *Adv Insect Physiol* 33:153–191
- Kleinhenz M, Bujok B, Fuchs S, Tautz J (2003) Hot bees in empty broodnest cells: heating from within. *J Exp Biol* 206:4217–4231
- Korb J (2003) Thermoregulation and ventilation of termite mounds. *Naturwissenschaften* 90:212–219
- Kronenberg F, Heller HC (1982) Colonial thermoregulation in honey bees (*Apis mellifera*). *J Comp Physiol A* 148:65–76
- Prange HD (1996) Evaporative cooling in insects. *J Insect Physiol* 42:493–499
- R Core Team (2013) R: a language and environment for statistical computing. R Foundation for Statistical Computing. <http://www.R-project.org>
- Sammataro D, Avitabile A (2011) Summer/fall management. In: *The beekeeper's handbook*, 4th edn. Cornell University Press, New York, pp 109–124
- Seeley TD, Morse RA (1976) The nest of the honey bee (*Apis mellifera* L.). *Insect Soc* 23:495–512
- Seeley TD (1989) The honey bee colony as a superorganism. *Amer Sci* 77:546–553
- Siegel AJ, Hui J, Johnson RN, Starks PT (2005) Honey bee workers as mobile insulating units. *Insect Soc* 52:242–246

- Simpson J (1961) Nest climate regulation in honey bee colonies: honey bees control their domestic environment by methods based on their habit of clustering together. *Science* 133:1327–1333
- Stabentheiner A, Pressl H, Papst T, Hrassnigg N, Crailsheim K (2003) Endothermic heat production in honeybee winter clusters. *J Exp Biol* 206:353–358
- Starks PT, Gilley DC (1999) Heat shielding: a novel method of colonial thermoregulation in honeybees. *Naturwissenschaften* 86:438–440
- Starks PT, Johnson RN, Siegel AJ, Decelle MM (2005) Heat shielding: a task for youngsters. *Behav Ecol* 16:128–132
- Tautz J (2008) Cultivated intelligence. In: *The buzz about bees: biology of a superorganism*, 1st edn. Springer-Verlag, Berlin, pp 205–232
- Tautz J, Maier S, Groh C, Rossler W, Brockmann A (2003) Behavioral performance in adult honey bees is influenced by the temperature experienced during their pupal development. *Proc Natl Acad Sci U S A* 100:7343–7347
- von Frisch K (1967) Orientation when near the goal. In: *Dance language and orientation of bees*. Harvard University Press, Massachusetts, pp 465–523
- Wheeler WM (1910) *Ants: their structure, development and behavior*. New York, New York
- Wilson EO (1971) Social homeostasis and the superorganism. In: *The insect societies*. Harvard University Press, Massachusetts, pp 306–319
- Winston ML (1987) *The biology of the honey bee*. Harvard University Press, Massachusetts